

March 9, 2004

MAZUR SEMINAR

1. INTRODUCTION

The goal of these notes is to show that the modular jacobian $J_0(N)_{/\mathbf{Q}}$ has a nontrivial quotient of rank 0 for any prime N such that the genus of $X_0(N)$ is positive.

Recall the following theorem from James' talk:

Theorem 1.1. *Let $A_{/\mathbf{Q}}$ be an abelian variety with good reduction outside of N and purely toric reduction at N . Suppose moreover that $\mathcal{A}[p]$ is admissible for some $p \neq N$, where $\mathcal{A}_{/\mathbf{Z}}$ is the Néron model of A . Then $A_{/\mathbf{Q}}$ has rank 0.*

Our aim is to construct a nonzero isogeny factor $J_{\mathfrak{P}}$ of $J_0(N)$ satisfying the hypotheses of this theorem. Before we construct $J_{\mathfrak{P}}$ however, we will show that the above reduction properties of A are inherited by any isogeny factor, and will discuss the relationship between isogenies $A \sim A' \times A''$ (with A, A', A'' abelian varieties over a field K) and idempotents in the ring of endomorphisms $\text{End}^0(A)$ in the isogeny category of abelian varieties over K .

2. SEMI-ABELIAN REDUCTION

In order to apply Theorem 1.1 to the isogeny factor $J_{\mathfrak{P}}$ of $J_0(N)$ that we will construct, we must show that $J_{\mathfrak{P}}$ has good reduction outside N and purely toric reduction at N , and that $J_{\mathfrak{P}}[p]$ is admissible. We first show that all isogeny factors of $J_{\mathfrak{P}}$ inherit these reduction properties.

Let K be the fraction field of a discrete valuation ring \mathcal{O} with normalized valuation v and residue field k . Fix a separable closure K_s of K and a prime v_s of K_s lying over v , and denote by I the inertia group of v_s . Let A be an abelian variety over K , and denote by \mathcal{A} its Néron model over \mathcal{O} , and by \mathcal{A}^0 the connected component of the identity of \mathcal{A} . Recall that we say that A has *semi-abelian reduction* with respect to \mathcal{O} if \mathcal{A}^0 is an extension of an abelian variety by a torus.

The aim of this section is to prove the following proposition:

Proposition 2.1. *Let the notation be as above. Suppose moreover that A is isogenous to the product $A' \times A''$ of abelian varieties A' and A'' over K . Then if A has good (resp. semi-abelian) reduction, so do A' and A'' .*

Proof. We will only prove the case of semi-abelian reduction as the other cases are easier. Choose a prime $\ell \neq \text{char } k$. Let $V_{\ell}(A) := T_{\ell}(A) \otimes \mathbf{Q}$ denote the rational ℓ -adic Tate module of A . First observe that

$$V_{\ell}(A) = V_{\ell}(A' \times A'') = V_{\ell}(A') \times V_{\ell}(A'').$$

Grothendieck's criterion for semi-abelian reduction (cf. [1], theorem 6, page 184) asserts that A has semi-abelian reduction if and only if there exists a subspace V of $V_{\ell}(A)$ stable under the inertia group I such that I acts trivially on both V and the quotient $V_{\ell}(A)/V$. (cf. [1, Theorem 6, p. 184]) Note that V can (and will) be taken to be the maximal subspace of $V_{\ell}(A)$ on which the action of I is trivial.

Let V', V'' be the maximal subspaces of $V_{\ell}(A')$ and $V_{\ell}(A'')$ respectively, on which I operates trivially. Obviously $V' \times V'' = V$. Moreover, as $(V_{\ell}(A')/V') \times (V_{\ell}(A'')/V'') = V_{\ell}(A)/V$, we conclude that I acts trivially on $V_{\ell}(A')/V'$ and on $V_{\ell}(A'')/V''$. This finishes the proof. \blacksquare

Remark 2.2. Let us prove that if A has toric reduction, then so do A' and A'' . Assume that A has toric reduction. First note that since A and $A' \times A''$ are isogenous, they are both semi-abelian as their rational Tate modules are isomorphic. Denote by ϕ_K an isogeny $A \rightarrow A' \times A''$. Then there exists an isogeny $\psi_K : A' \times A'' \rightarrow A$ and a nonzero integer n , such that $\phi_K \psi_K = n_{A' \times A''}$ and $\psi_K \phi_K = n_A$ [5, p. 169]. The Néron mapping property ensures that ϕ_K and ψ_K extend to morphisms ϕ and ψ between \mathcal{A} and the Néron model $\mathcal{A}' \times \mathcal{A}''$ of $A' \times A''$. We have $\text{Ner}(A' \times A'') = \mathcal{A}' \times \mathcal{A}''$ because of the Néron mapping property: if Z is a smooth \mathcal{O} -scheme, then

$$\begin{aligned} \text{Hom}_K(Z_K, A' \times A'') &= \text{Hom}_K(Z_K, A') \times \text{Hom}_K(Z_K, A'') \\ &= \text{Hom}_{\mathcal{O}}(Z, \mathcal{A}') \times \text{Hom}_{\mathcal{O}}(Z, \mathcal{A}'') = \text{Hom}_{\mathcal{O}}(Z, \mathcal{A}' \times \mathcal{A}''). \end{aligned}$$

We will first show that $\bar{\phi} : \bar{\mathcal{A}} \rightarrow \bar{\mathcal{A}}' \times \bar{\mathcal{A}}''$ is an isogeny. First note that $\bar{\psi}\bar{\phi} = \bar{\psi}\phi = [n]$. Since A is semi-abelian, $\bar{\mathcal{A}}^0$ is an extension of an abelian variety by a torus. As multiplication by a nonzero integer is surjective on abelian varieties and on tori, we conclude that $[n]$ is surjective on $\bar{\mathcal{A}}^0$, hence an isogeny. Since $[n] = \bar{\psi}\bar{\phi}$, and $\bar{\mathcal{A}}$ and $\bar{\mathcal{A}}' \times \bar{\mathcal{A}}''$ have the same dimension, we conclude that $\bar{\phi}$ is surjective on the identity components and has finite kernel, hence is an isogeny.

The isogeny between $\bar{\mathcal{A}}$ and $\bar{\mathcal{A}}' \times \bar{\mathcal{A}}''$ induces an isogeny between $\bar{\mathcal{A}}^0$ and $(\bar{\mathcal{A}}' \times \bar{\mathcal{A}}'')^0 = \bar{\mathcal{A}}'^0 \times \bar{\mathcal{A}}''^0$, so it remains to check that for connected commutative smooth algebraic groups over a field k , the property of being a torus is preserved under isogeny and under formation of direct factors. This is clear since by the structure theorem for commutative algebraic groups over \bar{k} such a group G is a torus if and only if G is affine and $\dim(V_\ell(G)) = \dim(G)$, where $\ell \neq \text{char } k$ is a prime.

3. IDEMPOTENTS INSIDE THE ENDOMORPHISM RING OF ABELIAN VARIETIES

In this section we remark upon some generalities concerning the relationship between idempotents in the ring $\text{End}^0(A)$ for an abelian variety A and isogenies $A \sim A' \times A''$ with A', A'' abelian varieties.

Proposition 3.1. *Let A be an abelian variety over a field K . There is a one-to-one correspondence between ordered pairs (A', A'') of abelian subvarieties of A such that $A' \times_K A'' \rightarrow A$ is an isogeny and idempotents in the ring $\text{End}^0(A)$; the operation $(A', A'') \mapsto (A'', A')$ corresponds to $e \mapsto 1 - e$.*

Proof. Suppose we are given a pair (A', A'') of abelian subvarieties of A such that the natural map $\phi : A' \times A'' \rightarrow A$ is an isogeny. Let ϕ^{-1} denote the inverse of ϕ inside $\text{Hom}^0(A, A' \times A'')$. Consider the composition

$$e : A \xrightarrow{\phi^{-1}} A' \times A'' \xrightarrow{\text{pr}} A' \xrightarrow{\iota} A' \times A'' \xrightarrow{\phi} A,$$

where ι is inclusion and pr is projection. Note that e is an idempotent in $\text{End}^0(A)$, and by definition of ϕ , it lifts the identity on A' .

Conversely, given an idempotent $e \in \text{End}^0(A)$, choose a nonzero integer n such that $ne \in \text{End}(A)$, and set A' to be the image of ne . Note that A' is independent of n , since multiplication by a nonzero integer is surjective on an abelian variety. The Poincaré Reducibility Theorem (*c.f.* [4]) guarantees the existence of a unique abelian subvariety A'' of A , such that $A' \times A'' \rightarrow A$ is an isogeny (the map being addition). One checks that the two processes are inverses of each other. ■

4. THE OPTIMAL QUOTIENT $J_{\mathfrak{P}}$ OF $J_0(N)$

In this section we will construct a nonzero optimal quotient $J_{\mathfrak{P}}$ of $J_0(N)$ such that the action of \mathbf{T} on $J_0(N)$ induces an action on $J_{\mathfrak{P}}$.

Fix N and let \mathbf{T} denote the Hecke ring. Let $\mathcal{I} \subseteq \mathbf{T}$ be the Eisenstein ideal let $\mathfrak{P} \supseteq \mathcal{I}$ be a prime of residual characteristic $p \neq N$ (we have seen that such a \mathfrak{P} exists). Observe that $\mathfrak{P}_p := \mathfrak{P}(\mathbf{T} \otimes \mathbf{Z}_p)$ is a prime of $T \otimes \mathbf{Z}_p$.

Consider the subring $\mathbf{T} \otimes_{\mathbf{Z}} \mathbf{Q} \subseteq \text{End}^0(J_0(N))$ (we have seen that in fact equality holds, but we do not use this). By Proposition 3.1, the decomposition of $\mathbf{T} \otimes \mathbf{Q}$ into a product of fields gives us the isogeny decomposition of $J_0(N)$ as a product of \mathbf{Q} -simple abelian subvarieties:

$$(1) \quad J_0(N) \sim \prod_{\mathfrak{q} \in \text{MinSpec}(\mathbf{T})} J_{\mathfrak{q}}.$$

We put

$$\tilde{J}_{\mathfrak{P}} = \prod_{\substack{\mathfrak{q} \in \text{MinSpec}(\mathbf{T}) \\ \mathfrak{q} \subseteq \mathfrak{P}}} J_{\mathfrak{q}}.$$

Definition 4.1. Let A, A' , and A'' be abelian varieties and suppose that A' and A'' are quotients of A . We say that A' and A'' are *isogenous as quotients* of A if there exists an isomorphism $\phi : A' \rightarrow A''$ in the isogeny category commuting (in that category) with the quotient maps $A \rightarrow A'$ and $A \rightarrow A''$.

Claim 4.2. *Let $A \rightarrow A''$ be a surjective map of abelian varieties over a field K . Then there is a unique quotient abelian variety A_{opt} , isogenous to A'' as a quotient of A , that is the quotient of A by an abelian subvariety.*

Proof. We treat existence first. By the Poincaré Reducibility Theorem over K , we see that A is isogenous to a product of simple abelian subvarieties, say $A \leftarrow \prod_{i=1}^n A_i$, and since A'' is a quotient of A we have (renumbering if necessary) an isogeny $A'' \sim \prod_{i=1}^k A_i$, as quotients of A , for some $k \leq n$. Define $A' = \prod_{k < i \leq n} A_i$. Then we have a morphism

$$A' \xhookrightarrow{\iota} \prod_{i=1}^n A_i \xrightarrow{\varphi} A,$$

where φ is an isogeny. Observe that $\varphi \circ \iota(A')$ is an abelian subvariety of A . We set

$$A_{\text{opt}} = A/\varphi \circ \iota(A').$$

It is clear that A_{opt} is the quotient of A by an abelian subvariety and it follows (again from Poincaré Reducibility) that A_{opt} is isogenous to A'' as a quotient of A .

We claim that A_{opt} has the following universal property: if $q'' : A \rightarrow \tilde{A}$ is any quotient isogenous to the quotient $q : A \rightarrow A_{\text{opt}}$ in the sense of Definition (4.1) then q'' factors uniquely through q in the isogeny category. Indeed, if \tilde{A} and A_{opt} are isogenous as quotients, there exists an isomorphism (In the isogeny category) $\phi : A_{\text{opt}} \rightarrow \tilde{A}$. By composing ϕ with multiplication by a large enough integer, we obtain *honest* maps $n \circ q'' : A \rightarrow \tilde{A}$ and $\psi = n \circ \phi : A_{\text{opt}} \rightarrow \tilde{A}$ such that $\psi \circ q = n \circ q''$. Hence $n \circ q''(\ker q) = 0$. But multiplication by n is an isogeny, and hence has finite kernel, while $q''(\ker q)$ is connected (as $\ker q$ is an abelian subvariety of A), so that $q''(\ker q) = 0$.

It follows that the map $A \rightarrow A''$ factors uniquely through A_{opt} . From this universal property of A_{opt} , we see at once that A_{opt} is unique up to unique isomorphism. \blacksquare

Applying the above claim to $A = J_0(N)$ and the isogeny factor $\tilde{J}_{\mathfrak{p}}$, we obtain an optimal quotient $J_{\mathfrak{p}}$ of $J_0(N)$. We claim that \mathbf{T} acts on $J_{\mathfrak{p}}$. This will follow from the following more general theorem:

Theorem 4.3. *Let $\pi : A \rightarrow A'$ be a surjective map of abelian varieties over a field K of characteristic 0 having an abelian variety kernel B and let $T \in \text{End}(A)$. Assume there exists $T' \in \text{End}(A')^0$ such that the following diagram commutes in the isogeny category:*

$$(2) \quad \begin{array}{ccc} A & \xrightarrow{\quad} & A' \\ & \searrow \pi & \\ T \downarrow & & \downarrow T' \\ A & \xrightarrow{\quad \pi} & A' \end{array}$$

Then $T' \in \text{End}(A')$.

Proof. We have $T_0 := nT' \in \text{End}(A')$ for some nonzero integer n , and by the universal property of the quotient map $A' \rightarrow A'$ having kernel $A'[n]$, we have $T' \in \text{End}(A')$ if and only if T_0 kills $A'[n]$. Since $T_0\pi = nT'\pi = n\pi T = \pi Tn$ in the isogeny category, the genuine maps $T_0\pi$ and πTn agree so that $T_0\pi$ kills $A'[n]$ as πTn obviously does. To conclude that T_0 kills $A'[n]$, it therefore suffices to show that π is faithfully flat on n -torsion, for which it is enough to show surjectivity on \overline{K} -points (as finite K -groups are étale so $A[n] \xrightarrow{\pi} A'[n]$ is faithfully flat if and only if it is surjective on \overline{K} -points). But we have the commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & B(\overline{K}) & \longrightarrow & A(\overline{K}) & \longrightarrow & A'(\overline{K}) \longrightarrow 0 \\ & & n \downarrow & & n \downarrow & & n \downarrow \\ 0 & \longrightarrow & B(\overline{K}) & \longrightarrow & A(\overline{K}) & \longrightarrow & A'(\overline{K}) \longrightarrow 0 \end{array}$$

and since A, A', B are abelian varieties and K is of characteristic 0, the vertical maps are all surjective, so by the Snake lemma, the map on n -torsion $A[n] \xrightarrow{\pi} A'[n]$ is surjective as desired. This completes the proof. \blacksquare

Applying Theorem 4.3 to the optimal quotient $J_{\mathfrak{p}}$ of $J_0(N)$ shows that we have an action of \mathbf{T} on $J_{\mathfrak{p}}$.

5. ADMISSIBILITY

In Brian's talk, it was explained why $J_{\mathfrak{P}}[p]$ is an admissible group scheme over \mathbf{Z} . In this section we will recall the proof of this fact in more detail.

Since \mathbf{T} is a finite \mathbf{Z} -module, the ring $\mathbf{T} \otimes_{\mathbf{Z}} \mathbf{Z}_p$ is a finite \mathbf{Z}_p -module and hence is semi-local. Moreover, we have seen in §4 that \mathbf{T} acts on $J_{\mathfrak{P}}$ in a manner that respects the quotient map $J_0(N) \rightarrow J_{\mathfrak{P}}$ so we obtain an action of $\mathbf{T} \otimes_{\mathbf{Z}} \mathbf{Z}_p$ on the Tate modules $T_p(J_0(N))$ and $T_p(J_{\mathfrak{P}})$ as $G_{\mathbf{Q}} := \text{Gal}(\overline{\mathbf{Q}}/\mathbf{Q})$ -modules. By 8.7 and 8.15 of [3], there is a canonical isomorphism

$$(3) \quad \mathbf{T} \otimes_{\mathbf{Z}} \mathbf{Z}_p \simeq \prod_{\mathfrak{m} \in \text{MaxSpec}(\mathbf{T} \otimes_{\mathbf{Z}} \mathbf{Z}_p)} (\mathbf{T} \otimes_{\mathbf{Z}} \mathbf{Z}_p)_{\mathfrak{m}}.$$

Claim 5.1. *The action of $\mathbf{T} \otimes_{\mathbf{Z}} \mathbf{Z}_p$ on $T_p(J_{\mathfrak{P}})$ factors through $\mathbf{T}_{\mathfrak{P}} := (\mathbf{T} \otimes_{\mathbf{Z}} \mathbf{Z}_p)_{\mathfrak{P}}$, so the induced action of $\mathbf{T} \otimes_{\mathbf{Z}} \mathbf{Z}_p$ on $J_{\mathfrak{P}}[p]$ is through the quotient $\mathbf{T}_{\mathfrak{P}}$.*

Proof. First note that the functoriality of the idempotent decomposition of $\mathbf{T} \otimes \mathbf{Q}$ implies that

$$V_p(J_{\mathfrak{P}}) = \prod_{\substack{\mathfrak{q} \in \text{Spec}(\mathbf{T} \otimes \mathbf{Q}) \\ \mathfrak{q} \subset \mathfrak{P}}} V_p(J_{\mathfrak{q}}) = \prod_{\substack{\mathfrak{q} \in \text{Spec}(\mathbf{T} \otimes \mathbf{Q}) \\ \mathfrak{q} \subset \mathfrak{P}}} V_p(J)_{\mathfrak{q}},$$

where the objects $V_p(J)_{\mathfrak{q}}$ denote the localizations of the $\mathbf{T} \otimes \mathbf{Q}$ -module $V_p(J)$ at primes \mathfrak{q} . (We will use the same letter \mathfrak{q} to denote both a prime ideal of $\mathbf{T} \otimes \mathbf{Q}$ and its inverse image in \mathbf{T}). From this we see that the action of \mathbf{T} on $V_p(J_{\mathfrak{P}})$ factors through $\prod_{\mathfrak{q} \subset \mathfrak{P}} (\mathbf{T} \otimes \mathbf{Q})_{\mathfrak{q}}$ and since every element of $\mathbf{T} - \mathfrak{P}$ is mapped to a unit of $\prod_{\mathfrak{q} \subset \mathfrak{P}} (\mathbf{T} \otimes \mathbf{Q})_{\mathfrak{q}}$ (since $\mathfrak{q} \subset \mathfrak{P}$), the map $\mathbf{T} \rightarrow \prod_{\mathfrak{q} \subset \mathfrak{P}} (\mathbf{T} \otimes \mathbf{Q})_{\mathfrak{q}}$ factors through the localization $\mathbf{T}_{\mathfrak{P}}$. The claim now follows after tensoring with \mathbf{Z}_p and noting that $T_p(J_{\mathfrak{P}}) \subseteq V_p(J_{\mathfrak{P}})$. ■

We now prove:

Lemma 5.2. *Let $\mathcal{I} \subseteq \mathbf{T}$ be the Eisenstein ideal and $\mathfrak{P} \supset \mathcal{I}$ be any prime of \mathbf{T} having residue characteristic $p \neq N$. Then \mathfrak{P}^r kills $J_{\mathfrak{P}}[p]$ for some $r > 0$.*

Proof. The action of $\mathbf{T} \otimes_{\mathbf{Z}} \mathbf{Z}_p$ on $J_{\mathfrak{P}}[p]$ factors through $\mathbf{T}_{\mathfrak{P}}/p\mathbf{T}_{\mathfrak{P}}$. As $\mathbf{T} \otimes_{\mathbf{Z}} \mathbf{Z}_p$ is a finite \mathbf{Z}_p -module, we see that $\mathbf{T}_{\mathfrak{P}}/p\mathbf{T}_{\mathfrak{P}}$ is a finite \mathbf{F}_p -module, and consequently is a finite local ring. It follows that its maximal ideal is nilpotent. Thus, for some $r > 0$ we have $\mathfrak{P}^r \subseteq p\mathbf{T}_{\mathfrak{P}}$, so $\mathfrak{P}^r J_{\mathfrak{P}}[p] = 0$ as claimed. ■

It follows from Lemma 5.2 that we have a filtration of $\mathbf{T}_{\mathfrak{P}}[G_{\mathbf{Q}}]$ modules

$$(4) \quad J_{\mathfrak{P}}[p] \supseteq \mathfrak{P} J_{\mathfrak{P}}[p] \supseteq \mathfrak{P}^2 J_{\mathfrak{P}}[p] \supseteq \dots \supseteq \mathfrak{P}^r J_{\mathfrak{P}}[p] = 0.$$

Observe that each quotient $\mathfrak{P}^i J_{\mathfrak{P}}[p] / \mathfrak{P}^{i+1} J_{\mathfrak{P}}[p]$ is killed by \mathfrak{P} . We will now show that this implies that the quotients are all admissible. Recall from [2, Corollary 1.6] that admissibility of a group scheme over \mathbf{Z} can be checked on the corresponding Galois module of $\overline{\mathbf{Q}}$ points. It then follows that $J_{\mathfrak{P}}[p]$ is admissible, by general results on Jordan–Hölder series.

Lemma 5.3. *Let G be a finite discrete $G_{\mathbf{Q}}$ -module of p -power order on which \mathbf{T} acts and let $\mathfrak{P} \supseteq \mathcal{I}$ be a prime of \mathbf{T} having residual characteristic $p \neq N$ and containing the Eisenstein ideal. Assume that for any prime $\ell \nmid Np$ the inertia group I_{ℓ} at ℓ acts trivially on G , and hence that we have an action of $\text{Frob}_{\ell} \in \text{Gal}(\overline{\mathbf{F}}_{\ell}/\mathbf{F}_{\ell})$ on G . Suppose that the Eichler–Shimura relation*

$$\text{Frob}_{\ell}^2 - T_{\ell} \text{Frob}_{\ell} + \ell = 0$$

holds on G for all such ℓ . If \mathfrak{P} kills G then G has a filtration by admissible closed subgroups with successive quotients $\mathbf{Z}/p\mathbf{Z}$ or μ_p .

Proof. Let Γ be the discrete finite quotient of $\text{Gal}(\overline{\mathbf{Q}}/\mathbf{Q})$ through which the Galois action on G factors. By hypothesis, for $\ell \nmid Np$ we have that Frob_{ℓ} acts on G and

$$(5) \quad \text{Frob}_{\ell}^2 - T_{\ell} \text{Frob}_{\ell} + \ell = 0$$

on G . But as $\mathfrak{P} \supseteq \mathcal{I}$ kills G and $T_{\ell} \equiv \ell + 1 \pmod{\mathcal{I}}$, we see that Frob_{ℓ} acts on G with eigenvalues contained in $\{1, \ell\}$. Now any $\sigma \in \text{Gal}(\overline{\mathbf{Q}}/\mathbf{Q})$ also acts on $G^{\vee} = \text{Hom}(G, \mu_p)$ via $f^{\sigma}(g) = \sigma f(g^{\sigma^{-1}})$. Observe that if g is an eigenvector of $\sigma = \text{Frob}_{\ell}$ with eigenvalue ℓ then $f^{\sigma}(g) = \sigma f(\ell^{-1}g) = \ell f(\ell^{-1}g) = f(g)$, so that $f(g)$ is an eigenvector

with eigenvalue 1. Similarly, if g has eigenvalue 1 then $f(g)$ has eigenvalue ℓ . It follows that the eigenvalues $\{1, \ell\}$ of Frob_ℓ on $G \times G^\vee$ occur with the same multiplicity. Thus, the characteristic polynomial of Frob_ℓ on $G \times G^\vee$ over \mathbf{F}_p is $(X - 1)^d(X - \ell)^d$ for some d (which is the dimension of G over \mathbf{F}_p).

Consider now the $\mathbf{F}_p[G_{\mathbf{Q}}]$ -module $(\mathbf{Z}/p\mathbf{Z})^d \times \mu_p^d$. By extending Γ if necessary (and still preserving the finiteness of Γ) we can regard $(\mathbf{Z}/p\mathbf{Z})^d \times \mu_p^d$ as a finite discrete Γ -module. Since Frob_ℓ acts on μ_p with eigenvalue ℓ , we see that the characteristic polynomial of Frob_ℓ on $(\mathbf{Z}/p\mathbf{Z})^d \times \mu_p^d$ is also $(X - \ell)^d(X - 1)^d$. Now by the Tchebotarev density theorem, every $\gamma \in \Gamma$ is the image of Frob_ℓ for some $\ell \neq N, p$. Thus, every $\gamma \in \Gamma$ has the same characteristic polynomial on the two $\mathbf{F}_p[\Gamma]$ -modules $(\mathbf{Z}/p\mathbf{Z})^d \times \mu_p^d$ and G . Applying the Brauer–Nesbitt theorem, we conclude that these Γ -modules have the same semisimplifications. Since $(\mathbf{Z}/p\mathbf{Z})^d \times \mu_p^d$ has a filtration as a Galois module with successive quotients isomorphic to $\mathbf{Z}/p\mathbf{Z}$ or μ_p , so does G . ■

Corollary 5.4. *Let N, p, \mathfrak{P} be as before. Then $J_{\mathfrak{P}}[p]$ is admissible.*

Proof. By [2, Corollary 1.6], we need only check that the Galois module $J_{\mathfrak{P}}[p]$ has a $\text{Gal}(\overline{\mathbf{Q}}/\mathbf{Q})$ -stable filtration by μ_p 's and $\mathbf{Z}/p\mathbf{Z}$'s. Using Proposition 2.1 and Remark 2.2, we see that $J_{\mathfrak{P}}$ has toric reduction at N and good reduction outside N , so the inertia group at any $\ell \nmid Np$ acts trivially on the Galois module $J_{\mathfrak{P}}[p](\overline{\mathbf{Q}})$. Moreover, the Eichler–Shimura relations hold on $J_{\mathfrak{P}}[p]$, as they do in $J_0(N)$ and hence on the isogeny factor $J_{\mathfrak{P}}$. We may therefore apply Lemma 5.3 to the situation discussed after the proof of Lemma 5.2 to conclude that $J_{\mathfrak{P}}[p]$ is admissible. ■

Corollary 5.5. *The nonzero isogeny factor $J_{\mathfrak{P}}$ of $J_0(N)$ has rank 0.*

Proof. By Proposition 2.1 and Remark 2.2, we know that $J_{\mathfrak{P}}$ has good reduction away from N and purely toric reduction at N . As $J_{\mathfrak{P}}[p]$ is admissible by Corollary 5.4, we see that $J_{\mathfrak{P}}$ satisfies the hypotheses of Theorem 1.1 and hence has rank 0. ■

REFERENCES

- [1] S. Bosch, W. Lutkebhöhmert, M. Raynaud, *Néron models*, Springer-Verlag Berlin Heidelberg 1990.
- [2] Mazur, B. Modular curves and the Eisenstein ideal. *Publ. math. I.H.E.S.*, **47**, (2), 1977. pp. 33–186.
- [3] Matsumura, H. *Commutative Ring Theory*, Cambridge University Press, 1986.
- [4] Milne, J. Abelian Varieties, in *Arithmetic Geometry*.
- [5] Mumford, D. *Abelian Varieties*, Oxford University Press, 1970.