

Counting rules useful in probability, 2.4 and 2.5

Grethe Hystad

September 4, 2013

Counting rules useful in probability, 2.4

Theorem

Fundamental Principle of Counting:

If the first task of an experiment can result in n_1 possible outcomes and for each such outcome, the second task can result in n_2 possible outcomes, then there are $n_1 n_2$ possible outcomes for the two tasks together.

Multiplication Principle


Example

Suppose two patients, patient 1 and patient 2, are randomly selected to participate in an experiment. They receive either drug A, drug B, or a placebo P. The total possible outcomes is $(2)(3) = 6$. The outcomes of the experiment can be written as ordered pairs: $(1, A)$, $(1, B)$, $(1, P)$, $(2, A)$, $(2, B)$, $(2, P)$.

Multiplication Principle, example continues

Example

We can also illustrate the outcomes with a tree-diagram:



Example 1

■ Example

A box contains 4 balls, of which 1 is green and 3 are blue. Suppose two balls are selected at random. Let A denote the event that the two balls have different colors. Find $P(A)$.

Example 1

■ Example

A box contains 4 balls, of which 1 is green and 3 are blue. Suppose two balls are selected at random. Let A denote the event that the two balls have different colors. Find $P(A)$.

■ Solution

- *Number the balls, 1, 2, 3, 4, where ball 1 is green and balls 2, 3, 4 are blue.*
- *Total possible outcomes are $4 \cdot 3 = 12$.*
- *Because the balls are randomly selected, each of the 12 outcomes has a probability of $\frac{1}{12}$.*
- $P(A) = P(\{(1, 2)\} \cup \{(1, 3)\} \cup \{(1, 4)\} \cup \{(2, 1)\} \cup \{(3, 1)\} \cup \{(4, 1)\}) = \frac{6}{12} = \frac{1}{2}$.

Example 2

■ Example

Suppose you want to choose a four digit pin code for your debit card. Each digit can take the numbers 0, 1, ..., 9. How many possible choices do you have?

Example 2

■ Example

Suppose you want to choose a four digit pin code for your debit card. Each digit can take the numbers 0, 1, ..., 9. How many possible choices do you have?

■ Solution

Since each of the 10 digits could be chosen for each position, there are

$$10 \times 10 \times 10 \times 10 = 10000$$

possible choices.

In general, if we have n objects and we select r of these with replacement, and order is important, then there are n^r possible outcomes.

Example 3

■ Example

7 applicants have applied for 7 different jobs. How many ways can the jobs be filled.

Example 3

Example

7 applicants have applied for 7 different jobs. How many ways can the jobs be filled.

Solution

- *Here the selection is without replacement because the same person does not fill more than one position.*
- *There are 7 choices for filling the first position, 6 for the second, ..., and 1 for the last position.*
- *By the Multiplication Principle, the positions can be filled in $7 \times 6 \times 5 \cdots \times 1 = 7! = 5040$ ways.*

Example 4

■ Example

7 applicants have applied for three different jobs. How many ways can the jobs be filled.

Example 4

■ Example

7 applicants have applied for three different jobs. How many ways can the jobs be filled.

■ Solution

- *Here the selection is without replacement because the same person does not fill more than one position.*
- *Can be filled in $7 \times 6 \times 5 = 210$ ways.*

Permutations

■ Definition

A permutation of a set of objects is an arrangement of those objects in a definite order without repetitions.

Permutations

■ Definition

A permutation of a set of objects is an arrangement of those objects in a definite order without repetitions.

- The number of permutations of a set with n distinct elements is $n! = n(n - 1)(n - 2) \cdots 2 \cdot 1$.

Permutations

■ Definition

A permutation of a set of objects is an arrangement of those objects in a definite order without repetitions.

- The number of permutations of a set with n distinct elements is $n! = n(n-1)(n-2) \cdots 2 \cdot 1$.
- The number of k permutations of a set of n distinct elements is $n(n-1) \cdots (n-k+1) = \frac{n!}{(n-k)!}$.

Permutations

■ Definition

A permutation of a set of objects is an arrangement of those objects in a definite order without repetitions.

- The number of permutations of a set with n distinct elements is $n! = n(n-1)(n-2) \cdots 2 \cdot 1$.
- The number of k permutations of a set of n distinct elements is $n(n-1) \cdots (n-k+1) = \frac{n!}{(n-k)!}$.

■ Example

The number of permutations of the set $\{1, 2, 3\}$ is $3! = 6$. They are 123 132 213 231 312 321.

Permutations

■ Example

The number of 2-permutations of the set $\{1, 2, 3\}$ is $\frac{3!}{(3-2)!} = 6$.
They are 12 21 13 31 23 32.

Permutations

Example

The number of 2-permutations of the set $\{1, 2, 3\}$ is $\frac{3!}{(3-2)!} = 6$.
They are 12 21 13 31 23 32.

Theorem

Permutations: The number of ordered arrangements or permutations P_k^n of k objects selected from n distinct objects ($k \leq n$) is given by

$$P_k^n = n(n-1) \cdots (n-k+1) = \frac{n!}{(n-k)!}.$$

In Example 3, $P_3^7 = \frac{7!}{(7-3)!} = \frac{7!}{4!} = 7 \times 6 \times 5 = 210$.

Combinations

Definition

A combination of a set of objects is a subset of those objects.

Theorem

Combinations: The number of distinct subsets or combinations of size k that can be selected from n distinct objects ($k \leq n$) is given by

$$C_k^n = \binom{n}{k} = \frac{n!}{k!(n-k)!}.$$

This is the same as the number of ways of partitioning n distinct objects into exactly two subsets containing k and $(n - k)$ elements, respectively.

Combinations

In R, the command for $\binom{n}{k}$ is: choose(n,k)

Example

There are $\binom{3}{2} = \frac{3!}{2!(3-2)!} = 3$ subsets of size 2 of the set $\{1, 2, 3\}$.
They are $\{1, 2\}$, $\{1, 3\}$, $\{2, 3\}$.

- If order matters, use permutations.
- If order does not matter, use combinations.

Combinations

- If a set S has n elements, there are

Combinations

- If a set S has n elements, there are
- $\binom{n}{0}$ subsets with 0 elements.

Combinations

- If a set S has n elements, there are
- $\binom{n}{0}$ subsets with 0 elements.
- $\binom{n}{1}$ subsets with 1 element.

Combinations

- If a set S has n elements, there are
- $\binom{n}{0}$ subsets with 0 elements.
- $\binom{n}{1}$ subsets with 1 element.
- $\binom{n}{2}$ subsets with 2 elements.

Combinations

- If a set S has n elements, there are
- $\binom{n}{0}$ subsets with 0 elements.
- $\binom{n}{1}$ subsets with 1 element.
- $\binom{n}{2}$ subsets with 2 elements.
- ..

Combinations

- If a set S has n elements, there are
- $\binom{n}{0}$ subsets with 0 elements.
- $\binom{n}{1}$ subsets with 1 element.
- $\binom{n}{2}$ subsets with 2 elements.
- ..
- ..

Combinations

- If a set S has n elements, there are
- $\binom{n}{0}$ subsets with 0 elements.
- $\binom{n}{1}$ subsets with 1 element.
- $\binom{n}{2}$ subsets with 2 elements.
- ..
- ..
- $\binom{n}{n}$ subsets with n elements.

Combinations

- If a set S has n elements, there are
- $\binom{n}{0}$ subsets with 0 elements.
- $\binom{n}{1}$ subsets with 1 element.
- $\binom{n}{2}$ subsets with 2 elements.
- ..
- ..
- $\binom{n}{n}$ subsets with n elements.
- The total number of possible subsets of S is $\binom{n}{0} + \binom{n}{1} + \cdots + \binom{n}{n}$.

Binomial formula

- Binomial Formula:

$$(1 + a)^n = \sum_{k=0}^{\infty} \binom{n}{k} a^k$$

which holds for all real n .

Binomial formula

- Binomial Formula:

$$(1 + a)^n = \sum_{k=0}^{\infty} \binom{n}{k} a^k$$

which holds for all real n .

- It follows that,

$$2^n = \sum_{k=0}^n \binom{n}{k}.$$

Binomial formula

- Binomial Formula:

$$(1 + a)^n = \sum_{k=0}^{\infty} \binom{n}{k} a^k$$

which holds for all real n .

- It follows that,

$$2^n = \sum_{k=0}^n \binom{n}{k}.$$

- Hence, if a set has n elements, it has 2^n different subsets.

Pascal's triangle

$$\binom{n}{k} = \binom{n-1}{k-1} + \binom{n-1}{k}.$$

Example

$$\binom{6}{3} = \binom{5}{2} + \binom{5}{3}.$$

Example

■ Example

In a Norway lottery, a player selects 6 numbers from 1 to 48. During the draw, 6 numbers are drawn at random and without replacement. To win the first prize, all 6 numbers must match those drawn in any order. How many winning numbers are possible?

Example

■ Example

In a Norway lottery, a player selects 6 numbers from 1 to 48. During the draw, 6 numbers are drawn at random and without replacement. To win the first prize, all 6 numbers must match those drawn in any order. How many winning numbers are possible?

■ Solution

The number of possible winning numbers is the combination of 6 objects selected from 48, that is,

$$\binom{48}{6} = \frac{48!}{6!(48-6)!} = 12271512.$$



Example

Since each of the 12271512 possible winning numbers are equally likely, the probability that a player who buys one ticket wins is $\frac{1}{12271512}$.

Example

A box contains 15 balls; 3 red, 5 blue and 7 green balls. 8 balls are selected from the box without replacement. What is the probability that the balls selected are 1 red , 4 blue and 3 green?

Solution

- The number of ways to select 8 balls from 15 is,
$$\binom{15}{8} = \frac{15!}{8!(15-8)!} = 6435.$$

Solution

- The number of ways to select 8 balls from 15 is,
$$\binom{15}{8} = \frac{15!}{8!(15-8)!} = 6435.$$
- The number of ways to select 1 red ball from 3 red is,
$$\binom{3}{1} = \frac{3!}{1!(3-1)!} = 3.$$

Solution

- The number of ways to select 8 balls from 15 is,
$$\binom{15}{8} = \frac{15!}{8!(15-8)!} = 6435.$$
- The number of ways to select 1 red ball from 3 red is,
$$\binom{3}{1} = \frac{3!}{1!(3-1)!} = 3.$$
- The number of ways to select 4 blue balls from 5 blue is,
$$\binom{5}{4} = \frac{5!}{4!(5-4)!} = 5.$$

Solution

- The number of ways to select 8 balls from 15 is,
$$\binom{15}{8} = \frac{15!}{8!(15-8)!} = 6435.$$
- The number of ways to select 1 red ball from 3 red is,
$$\binom{3}{1} = \frac{3!}{1!(3-1)!} = 3.$$
- The number of ways to select 4 blue balls from 5 blue is,
$$\binom{5}{4} = \frac{5!}{4!(5-4)!} = 5.$$
- The number of ways to select 3 green balls from 7 green is,
$$\binom{7}{3} = \frac{7!}{3!(7-3)!} = 35.$$

Solution

- The number of ways to select 8 balls from 15 is,
$$\binom{15}{8} = \frac{15!}{8!(15-8)!} = 6435.$$
- The number of ways to select 1 red ball from 3 red is,
$$\binom{3}{1} = \frac{3!}{1!(3-1)!} = 3.$$
- The number of ways to select 4 blue balls from 5 blue is,
$$\binom{5}{4} = \frac{5!}{4!(5-4)!} = 5.$$
- The number of ways to select 3 green balls from 7 green is,
$$\binom{7}{3} = \frac{7!}{3!(7-3)!} = 35.$$
- Then by the Fundamental Principle of Counting, the number of ways to select 1 red ball, 4 blue balls and 3 green is,
$$\binom{3}{1} \binom{5}{4} \binom{7}{3} = 3 \cdot 5 \cdot 35 = 525.$$

Solution

- The number of ways to select 8 balls from 15 is,
$$\binom{15}{8} = \frac{15!}{8!(15-8)!} = 6435.$$
- The number of ways to select 1 red ball from 3 red is,
$$\binom{3}{1} = \frac{3!}{1!(3-1)!} = 3.$$
- The number of ways to select 4 blue balls from 5 blue is,
$$\binom{5}{4} = \frac{5!}{4!(5-4)!} = 5.$$
- The number of ways to select 3 green balls from 7 green is,
$$\binom{7}{3} = \frac{7!}{3!(7-3)!} = 35.$$
- Then by the Fundamental Principle of Counting, the number of ways to select 1 red ball, 4 blue balls and 3 green is,
$$\binom{3}{1} \binom{5}{4} \binom{7}{3} = 3 \cdot 5 \cdot 35 = 525.$$
- The probability of selecting 1 red ball, 4 blue balls and 3 green balls is,
$$\frac{\binom{3}{1} \binom{5}{4} \binom{7}{3}}{\binom{15}{8}} = \frac{525}{6435} = \frac{105}{1287}.$$

Example

Example

A hand of seven cards is dealt from a deck of 52 cards.

- (A) How many hands are possible?
- (B) How many hands that contain exactly 2 diamonds are possible?
- (C) How many hands with 2 diamonds and 3 spades are possible?

Solution

Solution

- (A) *The number of seven-card hands is the number of choosing 7 elements from a set of 52 which is $\binom{52}{7}$.*

Solution

Solution

- (A) *The number of seven-card hands is the number of choosing 7 elements from a set of 52 which is $\binom{52}{7}$.*
- (B) *A hand with 2 diamonds can be formed as follows. Imagine,*
 - *deal 2 diamonds from the 13 diamonds.*
 - *deal 5 cards from the remaining 39 cards.*

The number of possible hands is,

$$\binom{13}{2} \binom{39}{5} = 44909046.$$

Solution

Solution

(C) A hand with 2 diamonds and 3 spades can be formed as follows.

- deal 2 diamonds from the 13 diamonds.
- deal 3 spades from the 13 spades.
- deal 2 cards from the remaining 26 cards.

The number of possible hands is,

$$\binom{13}{2} \binom{13}{3} \binom{26}{2} = 7250100.$$

Theorem

Partition: The number of ways of partitioning n distinct objects into k groups containing n_1, n_2, \dots, n_k objects, respectively, is

$$\frac{n!}{n_1!n_2! \cdots n_k!},$$

where

$$\sum_{i=1}^k n_i = n.$$

Example

■ Example

In how many ways can we permute the letters of the word MISSISSIPPI?

Example

■ Example

In how many ways can we permute the letters of the word MISSISSIPPI?

■ Solution

$$n = 11$$

$$n_1 = 1 \text{ M}$$

$$n_2 = 4 \text{ I's}$$

$$n_3 = 4 \text{ S's}$$

$$n_4 = 2 \text{ P's}$$

Hence, number of permutations is

$$\frac{11!}{1!4!4!2!} = 34650.$$



Example

■ Example

A coin is flipped 5 times, and the number of heads and tails is counted.

(A) How many flips have exactly 2 heads?

(B) How many flips have at most 2 heads?

(C) How many flips have at least 2 heads?

Example

Example

A coin is flipped 5 times, and the number of heads and tails is counted.

(A) How many flips have exactly 2 heads?

(B) How many flips have at most 2 heads?

(C) How many flips have at least 2 heads?

Solution

(A) If there are exactly 2 heads, there are exactly 3 tails, so the number of such flips is the number of permutations of 2 heads and 3 tails. That number is $\frac{5!}{2!3!} = 10$.

- (B) At most 2 heads means that one of the following is true:
 - no heads
 - exactly 1 head
 - exactly 2 heads
 - Hence $\frac{5!}{0!5!} + \frac{5!}{1!4!} + \frac{5!}{2!3!} = 1 + 5 + 10 = 16$ flips have at most 2 heads.

- (B) At most 2 heads means that one of the following is true:
 - no heads
 - exactly 1 head
 - exactly 2 heads
 - Hence $\frac{5!}{0!5!} + \frac{5!}{1!4!} + \frac{5!}{2!3!} = 1 + 5 + 10 = 16$ flips have at most 2 heads.
- (C) At least 2 heads means that one of the following is true:
 - exactly 2 heads
 - exactly 3 heads
 - exactly 4 heads
 - exactly 5 heads
 - Hence $\frac{5!}{2!3!} + \frac{5!}{3!2!} + \frac{5!}{4!1!} + \frac{5!}{5!0!} = 10 + 10 + 5 + 1 = 26$ flips have at least 2 heads.

Example

Example

A coin is flipped n times.

- Total number of outcomes is the number of sequences of length n which is 2^n .
- The number of sequences with k heads, is the number of subsets of size k of a set with n elements, which is $\binom{n}{k} = \frac{n!}{k!(n-k)!}$.
- The probability of k heads in a sequence of n coin tosses is

$$\frac{\binom{n}{k}}{2^n}.$$

Problem

Problem

A group of 9 people is divided into 3 groups of 3 people at random. Suppose that 3 of the people do not get along. What is the probability that the people are divided into groups such that all gets along?

Solution is given in class.

More counting rules useful in probability, chapter 2.5

Theorem

The number of ways of making k selections from n objects when selection is made with replacement and order is not important is

$$\binom{n + k - 1}{k}$$

Example

Example

Suppose in a lottery, a player selects 6 numbers from 1 to 48. During the draw, 6 numbers are drawn at random and with replacement. To win the first prize, all 6 numbers must match those drawn in any order. How many winning numbers are possible?

Solution

The number of possible winning numbers is

$$\binom{48 + 6 - 1}{6} = \frac{53 \times 52 \times 51 \times 50 \times 49 \times 48}{6 \times 5 \times 4 \times 3 \times 2 \times 1} = 22957480.$$

Summary

	Order is Important	Order is Not Important
With Replacement	n^k	$\binom{n+k-1}{k}$
Without Replacement	$P_k^n = \frac{n!}{(n-k)!}$	$C_k^n = \binom{n}{k}$

Birthday problem, chapter 2.4

Example

Suppose k people are in a room

(A) What is the probability that no two of them have the same birthday?

(B) How many people must be in the room for the probability that at least two of the k people have the same birthday to be greater than $\frac{1}{2}$.

Birthday problem, Solution, part (A)

Solution

(A) Since there are k people in the room and each has 365 possibilities for a birthday, there are 365^k possible outcomes. The number of possible outcomes for which no 2 have the same birthday is

$$(365)(364)\dots(365 - k + 1).$$

Thus, the probability, under the assumption that each outcome is equally likely, that no 2 people in the room have the same birthday is,

$$\frac{(365)(364)\dots(365 - k + 1)}{365^k}.$$

Birthday problem, Solution, part (B)

Solution

(B) The probability that at least two people of the k people share a birthday is

$$1 - \frac{(365)(364)\dots(365 - k + 1)}{365^k}.$$

Note that we can write

$$\frac{(365)(364)\dots(365 - k + 1)}{365^k} = \prod_{i=1}^k \left(1 - \frac{i-1}{365}\right).$$

Using R , we obtained that when $k = 23$, there is a 50% chance that at least two of the people in the room share a birthday. In a class of 38 students, there is a 86% chance that at least two people share a birthday.

Example

Suppose a box contains 30 balls, of which 12 is white and 18 is black. Pick 4 balls without replacement. The total possible outcomes are $\binom{30}{4}$. Let k be the number of white balls chosen. Then we have chosen k white balls out of 12 and $4 - k$ black balls out of 18. Then by the Fundamental Theorem of Counting, there are $\binom{12}{k} \cdot \binom{18}{4-k}$ possible outcomes. Thus, the probability of choosing k white balls is then

$$\frac{\binom{12}{k} \cdot \binom{18}{4-k}}{\binom{30}{4}}.$$

Example

```
> k <- -c(0 : 4)
> P <- -(choose(12, k) * choose(18, 4 - k))/(choose(30, 4))
> data.frame(k, P)
  k  P
1  0 0.11165846
2  1 0.35730706
3  2 0.36847291
4  3 0.14449918
5  4 0.01806240
```