

Symmetries of partial differential equations

Basic idea: get new solutions from existing ones using a transformation.

Symmetries of partial differential equations

Basic idea: get new solutions from existing ones using a transformation.

For example, if $u(x, t)$ solves $u_t = u_{xx}$, so does $u(x - x_0, t - t_0)$. This is a *translation* symmetry.

Symmetries of partial differential equations

Basic idea: get new solutions from existing ones using a transformation.

For example, if $u(x, t)$ solves $u_t = u_{xx}$, so does $u(x - x_0, t - t_0)$. This is a *translation* symmetry.

Also, if $u(x, t)$ solves $u_t = u_{xx}$, so does $u(-x, t)$. This is a *reflection* symmetry. Note $u(x, -t)$ is not a solution, however!

Symmetry transformations

Given a function $u(x, t)$, transform both independent and dependent variables by a mapping

$$(x, t, u) \rightarrow (x', t', u')$$

or explicitly

$$x' = X(x, t, u), \quad t' = T(x, t, u), \quad u' = U(x, t, u).$$

Symmetry transformations

Given a function $u(x, t)$, transform both independent and dependent variables by a mapping

$$(x, t, u) \rightarrow (x', t', u')$$

or explicitly

$$x' = X(x, t, u), \quad t' = T(x, t, u), \quad u' = U(x, t, u).$$

A partial differential equation is said to have a *symmetry* if

$$u'(x, t) = U(x, t, u(X(x, t, u), T(x, t, u)))$$

is a solution, given that $u(x, t)$ is.

Example 1

Consider equation

$$u_t = uu_{xx}$$

and transformation

$$x' = -x, \quad t' = -t, \quad u' = -u.$$

Example 1

Consider equation

$$u_t = uu_{xx}$$

and transformation

$$x' = -x, \quad t' = -t, \quad u' = -u.$$

Given solution $u(x, t)$, check $u'(x, t) = -u(-x, -t)$ also works:

$$\begin{aligned} \partial_t \left(-u(-x, -t) \right) &= u_t(-x, -t), \quad \text{and} \\ \left(-u(-x, -t) \right) \partial_{xx} \left(-u(-x, -t) \right) &= u(-x, -t) u_{xx}(-x, -t). \end{aligned}$$

Example 2

Consider the nonlinear Schrödinger equation

$$iu_t + \frac{1}{2}u_{xx} + |u|^2u = 0, \quad u(x, t) : \mathbb{R}^2 \rightarrow \mathbb{C}.$$

Example 2

Consider the nonlinear Schrödinger equation

$$iu_t + \frac{1}{2}u_{xx} + |u|^2u = 0, \quad u(x, t) : \mathbb{R}^2 \rightarrow \mathbb{C}.$$

"Galilean symmetry", with parameter V

$$U(x, t, u) = \exp(-iV(x + Vt/2))u, \quad X = x + Vt, \quad T = t.$$

Example 2

Consider the nonlinear Schrödinger equation

$$iu_t + \frac{1}{2}u_{xx} + |u|^2u = 0, \quad u(x, t) : \mathbb{R}^2 \rightarrow \mathbb{C}.$$

“Galilean symmetry”, with parameter V

$$U(x, t, u) = \exp(-iV(x + Vt/2))u, \quad X = x + Vt, \quad T = t.$$

Check $w(x, t) = \exp(-iV(x + Vt/2))u(x + Vt, t)$ is a solution:

$$\begin{aligned} iw_t + \frac{1}{2}w_{xx} + |w|^2w &= \\ e^{-iV(x+Vt/2)} \left\{ i \left(\frac{-iV^2}{2}u + u_t + Vu_x \right) + \frac{1}{2} \left(-V^2u - 2iVu_x + u_{xx} \right) + |u|^2u \right\} \\ &= e^{-iV(x+Vt/2)} \left\{ u_t + \frac{1}{2}u_{xx} + |u|^2u \right\} = 0. \end{aligned}$$

Dilation symmetries

Special type of symmetry involves rescaling space and time

$$x' = \frac{x}{L}, \quad t' = \frac{t}{L^\beta},$$

where $L > 0$.

Dilation symmetries

Special type of symmetry involves rescaling space and time

$$x' = \frac{x}{L}, \quad t' = \frac{t}{L^\beta},$$

where $L > 0$.

Mapping $u(x, t)$ to $u(x/L, t/L^\beta)$ is called a **dilation symmetry**.

Dilation symmetries

Special type of symmetry involves rescaling space and time

$$x' = \frac{x}{L}, \quad t' = \frac{t}{L^\beta},$$

where $L > 0$.

Mapping $u(x, t)$ to $u(x/L, t/L^\beta)$ is called a **dilation symmetry**.

Example: consider transport equation $u_t + cu_x = 0$.

Dilation symmetries

Special type of symmetry involves rescaling space and time

$$x' = \frac{x}{L}, \quad t' = \frac{t}{L^\beta},$$

where $L > 0$.

Mapping $u(x, t)$ to $u(x/L, t/L^\beta)$ is called a **dilation symmetry**.

Example: consider transport equation $u_t + cu_x = 0$.

Given solution $u(x, t)$, for which β is $u(x/L, t/L^\beta)$ also a solution?

Dilation symmetries

Special type of symmetry involves rescaling space and time

$$x' = \frac{x}{L}, \quad t' = \frac{t}{L^\beta},$$

where $L > 0$.

Mapping $u(x, t)$ to $u(x/L, t/L^\beta)$ is called a **dilation symmetry**.

Example: consider transport equation $u_t + cu_x = 0$.

Given solution $u(x, t)$, for which β is $u(x/L, t/L^\beta)$ also a solution?

$$\frac{u_t(x/L, t/L^\beta)}{L^\beta} + c \frac{u_x(x/L, t/L^\beta)}{L} = 0,$$

therefore need $\beta = 1$.

For example, given solution $u = \sin(x - ct)$, can construct a new solution $u = \sin([x - ct]/L)$.

Dilation symmetries, example 2

Consider symmetries $(x, y) \rightarrow (x/L, y/L^\beta)$ for

$$u_x + y^2 u_y = 0,$$

Dilation symmetries, example 2

Consider symmetries $(x, y) \rightarrow (x/L, y/L^\beta)$ for

$$u_x + y^2 u_y = 0,$$

Substituting $u(x/L, y/L^\beta)$ in for u ,

$$L^{-1} u_x(x/L, y/L^\beta) + L^{-\beta} y^2 u_y(x/L, y/L^\beta) = 0.$$

Dilation symmetries, example 2

Consider symmetries $(x, y) \rightarrow (x/L, y/L^\beta)$ for

$$u_x + y^2 u_y = 0,$$

Substituting $u(x/L, y/L^\beta)$ in for u ,

$$L^{-1} u_x(x/L, y/L^\beta) + L^{-\beta} y^2 u_y(x/L, y/L^\beta) = 0.$$

Caution! This is not the same as original equation since derivatives are evaluated where y is replaced with y/L^β , but the y^2 coefficient is not.

Dilation symmetries, example 2

Consider symmetries $(x, y) \rightarrow (x/L, y/L^\beta)$ for

$$u_x + y^2 u_y = 0,$$

Substituting $u(x/L, y/L^\beta)$ in for u ,

$$L^{-1} u_x(x/L, y/L^\beta) + L^{-\beta} y^2 u_y(x/L, y/L^\beta) = 0.$$

Caution! This is not the same as original equation since derivatives are evaluated where y is replaced with y/L^β , but the y^2 coefficient is not.

Remedy: rewrite as

$$L^{-1} u_x(x/L, y/L^\beta) + L^\beta (y/L^\beta)^2 u_y(x/L, y/L^\beta) = 0.$$

which means $\beta = -1$.

More general symmetry rescales dependent variable also:

$$(x, t, u) \rightarrow (x/L, t/L^\beta, u/L^\gamma).$$

More general symmetry rescales dependent variable also:

$$(x, t, u) \rightarrow (x/L, t/L^\beta, u/L^\gamma).$$

Exponents β, γ found by substituting $L^{-\gamma} u(x/L, t/L^\beta)$.

More general symmetry rescales dependent variable also:

$$(x, t, u) \rightarrow (x/L, t/L^\beta, u/L^\gamma).$$

Exponents β, γ found by substituting $L^{-\gamma} u(x/L, t/L^\beta)$.

Example: consider nonlinear convection diffusion equation

$$u_t = uu_{xx} - u_x.$$

Dilation symmetries, cont.

More general symmetry rescales dependent variable also:

$$(x, t, u) \rightarrow (x/L, t/L^\beta, u/L^\gamma).$$

Exponents β, γ found by substituting $L^{-\gamma} u(x/L, t/L^\beta)$.

Example: consider nonlinear convection diffusion equation

$$u_t = uu_{xx} - u_x.$$

Substitute $L^{-\gamma} u(x/L, t/L^\beta)$ for u ,

$$L^{-\gamma-\beta} u_t = L^{-2\gamma-2} uu_{xx} - L^{-\gamma-1} u_x, \quad \text{evaluated at } (x/L, t/L^\beta).$$

More general symmetry rescales dependent variable also:

$$(x, t, u) \rightarrow (x/L, t/L^\beta, u/L^\gamma).$$

Exponents β, γ found by substituting $L^{-\gamma} u(x/L, t/L^\beta)$.

Example: consider nonlinear convection diffusion equation

$$u_t = uu_{xx} - u_x.$$

Substitute $L^{-\gamma} u(x/L, t/L^\beta)$ for u ,

$$L^{-\gamma-\beta} u_t = L^{-2\gamma-2} uu_{xx} - L^{-\gamma-1} u_x, \quad \text{evaluated at } (x/L, t/L^\beta).$$

It follows $\gamma + \beta = 2\gamma + 2 = \gamma + 1$, so that $\beta = 1$ and $\gamma = -1$.

Suppose a PDE has a symmetry transformation $(x, t) \rightarrow (x/L, t/L^\beta)$.

Similarity solutions

Suppose a PDE has a symmetry transformation $(x, t) \rightarrow (x/L, t/L^\beta)$.

A solution $u = f(\eta)$ is called a **similarity solution**, where $\eta = x/t^{1/\beta}$ is called the **similarity variable**.

Similarity solutions

Suppose a PDE has a symmetry transformation $(x, t) \rightarrow (x/L, t/L^\beta)$.

A solution $u = f(\eta)$ is called a **similarity solution**, where $\eta = x/t^{1/\beta}$ is called the **similarity variable**.

Practical benefit: reduces problem to ODE for $f(\eta)$.

Similarity solutions

Suppose a PDE has a symmetry transformation $(x, t) \rightarrow (x/L, t/L^\beta)$.

A solution $u = f(\eta)$ is called a **similarity solution**, where $\eta = x/t^{1/\beta}$ is called the **similarity variable**.

Practical benefit: reduces problem to ODE for $f(\eta)$.

Similarity solutions are often physically important since they are scale invariant.

Invariance of similarity solutions

Fact: similarity solutions are the only ones which remain the same under symmetry transformation.

Invariance of similarity solutions

Fact: similarity solutions are the only ones which remain the same under symmetry transformation.

If $u(x, t) = f(\eta)$ is a similarity solution,

$$u(x/L, t/L^\beta) = f\left(\frac{x/L}{(t/L^\beta)^{1/\beta}}\right) = f\left(\frac{x}{t^{1/\beta}}\right) = u(x, t).$$

Invariance of similarity solutions

Fact: similarity solutions are the only ones which remain the same under symmetry transformation.

If $u(x, t) = f(\eta)$ is a similarity solution,

$$u(x/L, t/L^\beta) = f\left(\frac{x/L}{(t/L^\beta)^{1/\beta}}\right) = f\left(\frac{x}{t^{1/\beta}}\right) = u(x, t).$$

Conversely, if $u(x, t)$ is invariant under symmetry transformation, can be written in terms of new variables $w(\eta, \xi)$, where

$$\eta = \frac{x}{t^{1/\beta}}, \quad \xi = xt.$$

Invariance of similarity solutions

Fact: similarity solutions are the only ones which remain the same under symmetry transformation.

If $u(x, t) = f(\eta)$ is a similarity solution,

$$u(x/L, t/L^\beta) = f\left(\frac{x/L}{(t/L^\beta)^{1/\beta}}\right) = f\left(\frac{x}{t^{1/\beta}}\right) = u(x, t).$$

Conversely, if $u(x, t)$ is invariant under symmetry transformation, can be written in terms of new variables $w(\eta, \xi)$, where

$$\eta = \frac{x}{t^{1/\beta}}, \quad \xi = xt.$$

The transformation in terms of the new variables is $(\eta, \xi) \rightarrow (\eta, \xi/L^{\beta+1})$, so that

$$w(\eta, \xi) = w(\eta, \xi/L^{\beta+1}).$$

Invariance of similarity solutions

Fact: similarity solutions are the only ones which remain the same under symmetry transformation.

If $u(x, t) = f(\eta)$ is a similarity solution,

$$u(x/L, t/L^\beta) = f\left(\frac{x/L}{(t/L^\beta)^{1/\beta}}\right) = f\left(\frac{x}{t^{1/\beta}}\right) = u(x, t).$$

Conversely, if $u(x, t)$ is invariant under symmetry transformation, can be written in terms of new variables $w(\eta, \xi)$, where

$$\eta = \frac{x}{t^{1/\beta}}, \quad \xi = xt.$$

The transformation in terms of the new variables is $(\eta, \xi) \rightarrow (\eta, \xi/L^{\beta+1})$, so that

$$w(\eta, \xi) = w(\eta, \xi/L^{\beta+1}).$$

Differentiating with respect to L and setting $L = 1$,

$$w_\xi(\eta, \xi/L^{\beta+1})\xi = 0,$$

which means $w = w(\eta)$.

Example

Consider the diffusion equation

$$u_t = Du_{xx}, \quad -\infty < x < \infty.$$

Example

Consider the diffusion equation

$$u_t = Du_{xx}, \quad -\infty < x < \infty.$$

Find $u(x/L, t/L^\beta)$ is also a solution provided $\beta = 2$.

Example

Consider the diffusion equation

$$u_t = Du_{xx}, \quad -\infty < x < \infty.$$

Find $u(x/L, t/L^\beta)$ is also a solution provided $\beta = 2$.

Seek similarity solution $u = f(\eta)$ where $\eta = x/\sqrt{t}$. Substitution into equation gives

$$Df''(\eta) + \frac{\eta}{2}f'(\eta) = 0.$$

Example

Consider the diffusion equation

$$u_t = Du_{xx}, \quad -\infty < x < \infty.$$

Find $u(x/L, t/L^\beta)$ is also a solution provided $\beta = 2$.

Seek similarity solution $u = f(\eta)$ where $\eta = x/\sqrt{t}$. Substitution into equation gives

$$Df''(\eta) + \frac{\eta}{2}f'(\eta) = 0.$$

Separation of variables gives

$$f'(\eta) = Ce^{-\eta^2/4D},$$

therefore

$$f(\eta) = C_1 \operatorname{erf}(\eta/4D) + C_2, \quad \operatorname{erf}(x) \equiv \frac{2}{\sqrt{\pi}} \int_0^x e^{-y^2} dy.$$

More general similarity solutions

For the more general dilation symmetry,

$$(x, t, u) \rightarrow (x/L, t/L^\beta, u/L^\gamma).$$

seek solutions of form

$$u = t^{-\gamma/\beta} f(\eta), \quad \eta = x/t^{1/\beta}$$

More general similarity solutions

For the more general dilation symmetry,

$$(x, t, u) \rightarrow (x/L, t/L^\beta, u/L^\gamma).$$

seek solutions of form

$$u = t^{-\gamma/\beta} f(\eta), \quad \eta = x/t^{1/\beta}$$

These solutions are invariant under the transformation:

$$\begin{aligned} L^{-\gamma} u(x/L, t/L^\beta) &= L^{-\gamma} (t/L^\beta)^{-\gamma/\beta} f\left(\frac{x/L}{(t/L^\beta)^{1/\beta}}\right) \\ &= t^{-\gamma/\beta} f\left(\frac{x}{t^{1/\beta}}\right) = u(x, t). \end{aligned}$$

Example

Consider the nonlinear diffusion equation

$$u_t = uu_{xx} - u^3.$$

Example

Consider the nonlinear diffusion equation

$$u_t = uu_{xx} - u^3.$$

Plug in $L^{-\gamma}u(x/L, t/L^\beta)$,

$$\begin{aligned} \frac{1}{L^{\gamma+\beta}}u(x/L, t/L^\beta) &= \frac{1}{L^{2\gamma+2}}u(x/L, t/L^\beta)u_{xx}(x/L, t/L^\beta) \\ &\quad - \frac{1}{L^{3\gamma}}u(x/L, t/L^\beta)^3. \end{aligned}$$

Example

Consider the nonlinear diffusion equation

$$u_t = uu_{xx} - u^3.$$

Plug in $L^{-\gamma}u(x/L, t/L^\beta)$,

$$\begin{aligned} \frac{1}{L^{\gamma+\beta}}u(x/L, t/L^\beta) &= \frac{1}{L^{2\gamma+2}}u(x/L, t/L^\beta)u_{xx}(x/L, t/L^\beta) \\ &\quad - \frac{1}{L^{3\gamma}}u(x/L, t/L^\beta)^3. \end{aligned}$$

Need $\gamma + \beta = 2\gamma + 2 = 3\gamma$, so that $\gamma = 2$ and $\beta = 4$.

Example

Consider the nonlinear diffusion equation

$$u_t = uu_{xx} - u^3.$$

Plug in $L^{-\gamma}u(x/L, t/L^\beta)$,

$$\begin{aligned} \frac{1}{L^{\gamma+\beta}}u(x/L, t/L^\beta) &= \frac{1}{L^{2\gamma+2}}u(x/L, t/L^\beta)u_{xx}(x/L, t/L^\beta) \\ &\quad - \frac{1}{L^{3\gamma}}u(x/L, t/L^\beta)^3. \end{aligned}$$

Need $\gamma + \beta = 2\gamma + 2 = 3\gamma$, so that $\gamma = 2$ and $\beta = 4$.

Similarity solution will have the form

$$u = t^{-1/2}f(\eta), \quad \eta = x/t^{1/4}.$$

Example

Consider the nonlinear diffusion equation

$$u_t = uu_{xx} - u^3.$$

Plug in $L^{-\gamma}u(x/L, t/L^\beta)$,

$$\begin{aligned}\frac{1}{L^{\gamma+\beta}}u(x/L, t/L^\beta) &= \frac{1}{L^{2\gamma+2}}u(x/L, t/L^\beta)u_{xx}(x/L, t/L^\beta) \\ &\quad - \frac{1}{L^{3\gamma}}u(x/L, t/L^\beta)^3.\end{aligned}$$

Need $\gamma + \beta = 2\gamma + 2 = 3\gamma$, so that $\gamma = 2$ and $\beta = 4$.

Similarity solution will have the form

$$u = t^{-1/2}f(\eta), \quad \eta = x/t^{1/4}.$$

Plugging into equation gives

$$-\frac{1}{2}f - \frac{\eta}{4}f' = ff'' - f^3.$$

Need to find solution numerically!

Incomplete similarity

Sometimes scaling exponents are not uniquely determined; this situation is called **incomplete similarity**.

Incomplete similarity

Sometimes scaling exponents are not uniquely determined; this situation is called **incomplete similarity**.

Extra conditions are required to fix the value of the exponents.

Sometimes scaling exponents are not uniquely determined; this situation is called **incomplete similarity**.

Extra conditions are required to fix the value of the exponents.

Example: for diffusion equation $u_t = Du_{xx}$, inserting $L^{-\gamma}u(x/L, t/L^\beta)$ into the equation gives $L^{-\gamma-\beta}u_t = DL^{-\gamma-2}u_{xx}$. Thus $\beta = 2$ but γ is arbitrary.

Incomplete similarity

Sometimes scaling exponents are not uniquely determined; this situation is called **incomplete similarity**.

Extra conditions are required to fix the value of the exponents.

Example: for diffusion equation $u_t = Du_{xx}$, inserting $L^{-\gamma}u(x/L, t/L^\beta)$ into the equation gives $L^{-\gamma-\beta}u_t = DL^{-\gamma-2}u_{xx}$. Thus $\beta = 2$ but γ is arbitrary.

Already found solution with $\gamma = 0$,

$$f(\eta) = C_1 \operatorname{erf}(\eta/4D) + C_2,$$

but this does not decay at $\pm\infty$.

Incomplete similarity, cont.

Suppose we want $u(\pm\infty) = 0$. This implies conservation

$$\frac{d}{dt} \int_{-\infty}^{\infty} u(x, t) dx = 0.$$

Incomplete similarity, cont.

Suppose we want $u(\pm\infty) = 0$. This implies conservation

$$\frac{d}{dt} \int_{-\infty}^{\infty} u(x, t) dx = 0.$$

Inserting solution of form $u = t^\alpha f(x/t^{1/\beta})$ and changing variables

$$\frac{d}{dt} \int_{-\infty}^{\infty} t^{\alpha+1/\beta} f(\eta) d\eta = 0,$$

therefore $\alpha = -1/\beta = -1/2$.

Incomplete similarity, cont.

Suppose we want $u(\pm\infty) = 0$. This implies conservation

$$\frac{d}{dt} \int_{-\infty}^{\infty} u(x, t) dx = 0.$$

Inserting solution of form $u = t^\alpha f(x/t^{1/\beta})$ and changing variables

$$\frac{d}{dt} \int_{-\infty}^{\infty} t^{\alpha+1/\beta} f(\eta) d\eta = 0,$$

therefore $\alpha = -1/\beta = -1/2$.

Substituting $u = t^{-1/2} f(x/t^{1/2})$, get ODE

$$f'' + \frac{\eta}{2} f' + \frac{1}{2} f = f'' + \frac{1}{2} (\eta f)' = 0.$$

Using the condition $f(\pm\infty) = 0$, integration gives

$$f' = -\frac{1}{2} \eta f, \quad df/f = -\frac{1}{2} \eta d\eta, \quad f(\eta) = C e^{-\eta^2/4}.$$

Incomplete similarity, cont.

Suppose we want $u(\pm\infty) = 0$. This implies conservation

$$\frac{d}{dt} \int_{-\infty}^{\infty} u(x, t) dx = 0.$$

Inserting solution of form $u = t^\alpha f(x/t^{1/\beta})$ and changing variables

$$\frac{d}{dt} \int_{-\infty}^{\infty} t^{\alpha+1/\beta} f(\eta) d\eta = 0,$$

therefore $\alpha = -1/\beta = -1/2$.

Substituting $u = t^{-1/2} f(x/t^{1/2})$, get ODE

$$f'' + \frac{\eta}{2} f' + \frac{1}{2} f = f'' + \frac{1}{2} (\eta f)' = 0.$$

Using the condition $f(\pm\infty) = 0$, integration gives

$$f' = -\frac{1}{2} \eta f, \quad df/f = -\frac{1}{2} \eta d\eta, \quad f(\eta) = C e^{-\eta^2/4}.$$

With $C = 1/\sqrt{4\pi}$ get fundamental solution

$$u = (4\pi t)^{-1/2} e^{-x^2/4t}$$

Similarity solutions, Example 1

Consider nonlinear transport equation

$$u_t + uu_x = 0.$$

Symmetry of the form $u(x, t) \rightarrow u(x/L, t/L^\beta)$ leads to $\beta = 1$.

Similarity solutions, Example 1

Consider nonlinear transport equation

$$u_t + uu_x = 0.$$

Symmetry of the form $u(x, t) \rightarrow u(x/L, t/L^\beta)$ leads to $\beta = 1$.

Look for similarity solution $u = f(x/t)$. Plugging in,

$$-\frac{x}{t^2}f' + \frac{1}{t}ff' = 0$$

or

$$f'(f - \eta) = 0.$$

Similarity solutions, Example 1

Consider nonlinear transport equation

$$u_t + uu_x = 0.$$

Symmetry of the form $u(x, t) \rightarrow u(x/L, t/L^\beta)$ leads to $\beta = 1$.

Look for similarity solution $u = f(x/t)$. Plugging in,

$$-\frac{x}{t^2}f' + \frac{1}{t}ff' = 0$$

or

$$f'(f - \eta) = 0.$$

Either $f' = 0$, or $f = \eta$. Latter choice leads to solution

$$u(x, t) = \eta = \frac{x}{t}.$$

i.e. a rarefaction wave.

Similarity solutions, Example 2

A model for a convective thermal layer is

$$3yu_x - u_{yy} = 0, \quad u(0, y) = 1, \quad u(x, 0) = 0.$$

Similarity solutions, Example 2

A model for a convective thermal layer is

$$3yu_x - u_{yy} = 0, \quad u(0, y) = 1, \quad u(x, 0) = 0.$$

Treat x like the time variable, substitute $u(x/L^\beta, y/L)$

$$\frac{3y}{L^\beta} u_x(x/L^\beta, y/L) - \frac{1}{L^2} u_{yy}(x/L^\beta, y/L) = 0.$$

Similarity solutions, Example 2

A model for a convective thermal layer is

$$3yu_x - u_{yy} = 0, \quad u(0, y) = 1, \quad u(x, 0) = 0.$$

Treat x like the time variable, substitute $u(x/L^\beta, y/L)$

$$\frac{3y}{L^\beta} u_x(x/L^\beta, y/L) - \frac{1}{L^2} u_{yy}(x/L^\beta, y/L) = 0.$$

Note: to make this like original equation, y should be replaced by y/L everywhere.

Similarity solutions, Example 2

A model for a convective thermal layer is

$$3yu_x - u_{yy} = 0, \quad u(0, y) = 1, \quad u(x, 0) = 0.$$

Treat x like the time variable, substitute $u(x/L^\beta, y/L)$

$$\frac{3y}{L^\beta} u_x(x/L^\beta, y/L) - \frac{1}{L^2} u_{yy}(x/L^\beta, y/L) = 0.$$

Note: to make this like original equation, y should be replaced by y/L everywhere. Write as:

$$\frac{3(y/L)}{L^{\beta-1}} u_x(x/L^\beta, y/L) - \frac{1}{L^2} u_{yy}(x/L^\beta, y/L) = 0.$$

therefore $u(x/L^\beta, y/L)$ is still a solution if $\beta = 3$.

Similarity solutions, Example 2, cont.

Plug in similarity ansatz $u = f(\eta)$, $\eta = y/x^{1/3}$,

$$-\eta^2 f'(\eta) - f''(\eta) = 0, \quad f(0) = 0, \quad \lim_{\eta \rightarrow \infty} f(\eta) = 1.$$

Similarity solutions, Example 2, cont.

Plug in similarity ansatz $u = f(\eta)$, $\eta = y/x^{1/3}$,

$$-\eta^2 f'(\eta) - f''(\eta) = 0, \quad f(0) = 0, \quad \lim_{\eta \rightarrow \infty} f(\eta) = 1.$$

Separate variables $f''/f' = -\eta^2$ and integrate

$$f' = A e^{-\eta^3/3},$$

and integrate again

$$f = A \int_0^\eta e^{-s^3/3} ds + B.$$

Using the boundary conditions, $B = 0$ and

$$A = \left(\int_0^\infty e^{-s^3/3} ds \right)^{-1}.$$

Similarity solutions, Example 3

An unwinding fluid vortex is described by

$$v_t = \left(\frac{1}{r} [rv]_r \right)_r, \quad r > 0, \quad v(r, 0) = \frac{1}{r}.$$

Similarity solutions, Example 3

An unwinding fluid vortex is described by

$$v_t = \left(\frac{1}{r} [rv]_r \right)_r, \quad r > 0, \quad v(r, 0) = \frac{1}{r}.$$

Inserting $L^{-\gamma} v(r/L, t/L^\beta)$

$$\frac{1}{L^{\beta+\gamma}} v_t(r/L, t/L^\beta) = \frac{1}{L^{2+\gamma}} \left(\frac{1}{(r/L)} [(r/L)v(r/L, t/L^\beta)]_r \right)_r,$$

thus $\beta = 2$ but γ is undetermined.

Similarity solutions, Example 3

An unwinding fluid vortex is described by

$$v_t = \left(\frac{1}{r} [rv]_r \right)_r, \quad r > 0, \quad v(r, 0) = \frac{1}{r}.$$

Inserting $L^{-\gamma} v(r/L, t/L^\beta)$

$$\frac{1}{L^{\beta+\gamma}} v_t(r/L, t/L^\beta) = \frac{1}{L^{2+\gamma}} \left(\frac{1}{(r/L)} [(r/L)v(r/L, t/L^\beta)]_r \right)_r,$$

thus $\beta = 2$ but γ is undetermined.

To be compatible with initial condition, insert $L^{-\gamma} v(r/L, 0)$

$$L^{-\gamma} v(r/L, 0) = L^{-1} (r/L)^{-1}, \quad \text{therefore } \gamma = 1.$$

Similarity solutions, Example 3 cont.

Since

$$t^{-1/2}f(r/t^{1/2}) = \frac{1}{r}[\eta^{-1}f(r/t^{1/2})],$$

easier to look for solution of form

$$v = \frac{f(\eta)}{r}, \quad \eta = r^2/t, \quad \lim_{\eta \rightarrow \infty} f(\eta) = 1.$$

Similarity solutions, Example 3 cont.

Since

$$t^{-1/2}f(r/t^{1/2}) = \frac{1}{r}[\eta^{-1}f(r/t^{1/2})],$$

easier to look for solution of form

$$v = \frac{f(\eta)}{r}, \quad \eta = r^2/t, \quad \lim_{\eta \rightarrow \infty} f(\eta) = 1.$$

Substitution gives $f' + 4f'' = 0$ whose general solution is

$$f(r) = A + Be^{-\eta/4}.$$

Similarity solutions, Example 3 cont.

Since

$$t^{-1/2}f(r/t^{1/2}) = \frac{1}{r}[\eta^{-1}f(r/t^{1/2})],$$

easier to look for solution of form

$$v = \frac{f(\eta)}{r}, \quad \eta = r^2/t, \quad \lim_{\eta \rightarrow \infty} f(\eta) = 1.$$

Substitution gives $f' + 4f'' = 0$ whose general solution is

$$f(r) = A + Be^{-\eta/4}.$$

In original variables

$$v(r, t) = \frac{1}{r} \left(1 + B \exp \left(-\frac{r^2}{4t} \right) \right).$$

For bounded solution at origin, $B = -1$.

Similarity solutions, Example 4.

The porous medium equation is

$$u_t = (uu_x)_x.$$

If $L^{-\gamma}u(r/L, t/L^\beta)$ is a solution, need $\beta = 2 + \gamma$.

Similarity solutions, Example 4.

The porous medium equation is

$$u_t = (uu_x)_x.$$

If $L^{-\gamma}u(r/L, t/L^\beta)$ is a solution, need $\beta = 2 + \gamma$.

Equation is a conservation law with flux $J = -uu_x$, so that

$$\frac{d}{dt} \int_{-\infty}^{\infty} u(x, t) dx = 0, \quad \text{if } J(\pm\infty) = 0.$$

Similarity solutions, Example 4.

The porous medium equation is

$$u_t = (uu_x)_x.$$

If $L^{-\gamma}u(r/L, t/L^\beta)$ is a solution, need $\beta = 2 + \gamma$.

Equation is a conservation law with flux $J = -uu_x$, so that

$$\frac{d}{dt} \int_{-\infty}^{\infty} u(x, t) dx = 0, \quad \text{if } J(\pm\infty) = 0.$$

Inserting

$$u = t^{-\gamma/\beta} f(\eta), \quad \eta = x/t^{1/\beta},$$

then

$$\int_{-\infty}^{\infty} u(x, t) dx = t^{(1-\gamma)/\beta} \int_{-\infty}^{\infty} f(\eta) d\eta$$

which means that $\gamma = 1$, $\beta = 3$.

Similarity solutions, Example 4, cont.

Similarity solution $u = t^{-1/3}f(\eta)$, $\eta = x/t^{1/3}$ solves

$$-\frac{1}{3}(\eta f)' = (ff)'$$

Similarity solutions, Example 4, cont.

Similarity solution $u = t^{-1/3}f(\eta)$, $\eta = x/t^{1/3}$ solves

$$-\frac{1}{3}(\eta f)' = (ff)'$$

Integration twice (assuming $f'(0) = 0$ or $f(\pm\infty) = 0$) gives

$$f = B - \frac{\eta^2}{6}.$$

Similarity solutions, Example 4, cont.

Similarity solution $u = t^{-1/3}f(\eta)$, $\eta = x/t^{1/3}$ solves

$$-\frac{1}{3}(\eta f)' = (ff)'$$

Integration twice (assuming $f'(0) = 0$ or $f(\pm\infty) = 0$) gives

$$f = B - \frac{\eta^2}{6}.$$

Problem! f is not positive (unphysical), and $J(\pm\infty) \neq 0$.

Similarity solutions, Example 4, cont.

Similarity solution $u = t^{-1/3}f(\eta)$, $\eta = x/t^{1/3}$ solves

$$-\frac{1}{3}(\eta f)' = (ff)'$$

Integration twice (assuming $f'(0) = 0$ or $f(\pm\infty) = 0$) gives

$$f = B - \frac{\eta^2}{6}.$$

Problem! f is not positive (unphysical), and $J(\pm\infty) \neq 0$.

Solution: just take positive part, so that

$$u(x, t) = \begin{cases} t^{-1/3} \left(B - \frac{x^2}{6t^{2/3}} \right) & x^2 < 6Bt^{2/3}, \\ 0 & x^2 > 6Bt^{2/3}. \end{cases}$$

Observation: value of B determines total mass $\int u dx$, and this is constant in time.