

MATH 129  
FINAL EXAM REVIEW PACKET ANSWERS  
(Fall 2014)

1.  $\int_0^3 \left( 10\pi \sin\left(\frac{\pi}{3}t\right) + 30 \right) dt = 150$  people

2.  $\int_1^2 f(5-2x)dx = \frac{7}{2}$       Let  $u = 5 - 2x$  and change the endpoints.

3. a)  $\int \frac{t}{\sqrt{t+1}} dt = \frac{2}{3}(t+1)^{3/2} - 2(t+1)^{1/2} + c$       By the method of substitution with  $u = t + 1$ .

You can also use integration by parts with  $u = t$  and  $v' = (t+1)^{-1/2}$ . The result is equivalent, just written in a different form.  $\int \frac{t}{\sqrt{t+1}} dt = 2t(t+1)^{1/2} - \frac{4}{3}(t+1)^{3/2} + c$

b)  $\int \left( \frac{1}{z^2} + A \right)^2 dz = -\frac{1}{3z^3} - \frac{2A}{z} + A^2z + c$       Distribute.

c)  $\int 3^x e^x dx = \frac{1}{\ln(3e)} (3e)^x + c$       Rewrite  $3^x e^x = (3e)^x$ .

d)  $\int_0^1 \frac{\arctan y}{1+y^2} dy = \frac{\pi^2}{32}$       Substitution and change endpoints. Always simplify answer.

4. a)  $\int \frac{\ln(z^2+1)}{z^2} dz = -\frac{1}{z} \ln(z^2+1) + 2 \arctan z + c$       Let  $u = \ln(z^2+1)$  and  $v' = \frac{1}{z^2}$ .

b)  $\int x \arcsin(x^2) dx = \frac{1}{2} x^2 \arcsin(x^2) + \frac{1}{2} \sqrt{1-x^4} + c$       First make a substitution with  $w = x^2$ .  
Then let  $u = \arcsin(w)$  and  $v' = 1$ .

c)  $\int_0^1 x \cdot g''(x) dx = 3$       Let  $u = x$  and  $v' = g''$  for the first integration by parts.

5. a)  $\int \cos^2(3\theta+2) d\theta = \frac{1}{6} \cos(3\theta+2) \sin(3\theta+2) + \frac{1}{6} (3\theta+2) + c$       Let  $u = 3\theta + 2$  before using table formula # 18. If you use another approach, your answer will look different.

b)  $\int \frac{2}{4t^2 - 9} dt = \frac{1}{6} (\ln|2t - 3| - \ln|2t + 3|) + c$  Let  $u = 2t$  and factor the denominator before using table formula # 26. If you use another approach, your answer will look different.

c)  $\int \frac{dy}{\sqrt{y^2 + 8y + 15}} = \ln \left| (y + 4) + \sqrt{y^2 + 8y + 15} \right| + c$  Complete the square before using table formula # 29.

d)  $\int \frac{\sin(4\alpha)}{\cos^2(4\alpha) - \cos(4\alpha)} d\alpha = \frac{1}{4} (\ln|\cos(4\alpha)| - \ln|\cos(4\alpha) - 1|) + c$  Let  $u = \cos(4\alpha)$  and factor the denominator before using table formula # 26.

6. a)  $\int \frac{3y^3 + 5y - 1}{y^3 + y} dy = 3y - \ln|y| + \frac{1}{2} \ln|y^2 + 1| + 2 \arctan(y) + c$  First do long division, then use partial fractions  $\frac{A}{y} + \frac{By + C}{y^2 + 1}$ .

b)  $\int \frac{5z - 28}{6z^2 + z - 40} dz = \frac{4}{3} \ln|3z + 8| - \frac{1}{2} \ln|2z - 5| + c$  Use partial fractions  $\frac{A}{3z + 8} + \frac{B}{2z - 5}$ .

c)  $\int \frac{dx}{(5 - x^2)^{3/2}} = \frac{1}{5} \frac{x}{\sqrt{5 - x^2}} + c$  Let  $x = \sqrt{5} \sin(\theta)$ .

d)  $\int \frac{dt}{t^2 \sqrt{1 + t^2}} = -\frac{\sqrt{1 + t^2}}{t} + c$  Let  $t = \tan(\theta)$ .

7. a)  $\int_1^3 f'(x) e^{f(x)} dx = e^{11} - e^7$  Let  $u = f(x)$ .

b)  $\int_1^e \frac{f'(\ln x)}{x} dx = 2$  Let  $u = \ln x$ .

8.  $v = k(R^2 - r^2)$ .  $\frac{1}{R - 0} \int_0^R k(R^2 - r^2) dr = \frac{2}{3} kR^2$

9.  $E(X) = \int_{-\infty}^{\infty} xf(x) dx = \int_{-\infty}^0 xf(x) dx + \int_0^{\infty} xf(x) dx = 0 + \int_0^{\infty} x \frac{1}{7} e^{-x/7} dx = 7$

Use integration by parts or the table of integrals. Remember to use proper notation.

10.  $\int_0^1 e^{-t^2} dt \approx \frac{1}{2}e^{-1/16} + \frac{1}{2}e^{-9/16} \approx 0.75459794$

11. a), d), and e)

12. Trap, Right, Mid, and Left

13. a) The integral converges.  $\int_0^\infty \frac{1}{x^2+4} dx = \frac{\pi}{4}$  Use table formula # 24.

b) The integral converges.  $\int_1^\infty \frac{1}{2^x} dx = \frac{1}{2 \ln 2}$ .

c) The integral diverges.  $\int_0^1 \frac{e^x}{(e^x-1)^2} dx = \infty$  Let  $u = e^x - 1$ .

d) The integral converges.  $\int_{\pi/6}^{\pi/2} \frac{\sin x}{\sqrt{\cos x}} dx = 2\sqrt{\frac{\sqrt{3}}{2}}$  Let  $u = \cos x$

e) The integral diverges.  $\int_1^\infty \frac{dx}{(x-2)^3} = \int_1^2 \frac{dx}{(x-2)^3} + \int_2^\infty \frac{dx}{(x-2)^3}$ . The first integral diverges.

f) The integral converges.  $\int_5^\infty \frac{du}{u^2-16} = -\frac{1}{8} \ln\left(\frac{1}{9}\right) = \frac{1}{8} \ln(9)$  Use table formula # 26.

14.  $\int_m^\infty e^{-\left(\frac{x-m}{s}\right)^2} dx = \frac{s\sqrt{\pi}}{2}$  Let  $u = \frac{x-m}{s}$  and change the endpoints.

15. a) The integral converges. Rewrite as  $\int_0^\infty a \cdot f(x) dx = a \int_0^\infty f(x) dx$ .

b) The integral converges. Let  $u = ax$ .

c) The integral diverges. Rewrite as  $\int_0^\infty (a + f(x)) dx = \int_0^\infty a dx + \int_0^\infty f(x) dx$ .

d) The integral converges. Let  $u = a + x$ .

16. a) The integral converges. By comparison with  $\int_2^{\infty} \frac{d\theta}{\theta^{3/2}}$ .
- b) The integral converges. By comparison with  $\int_1^{\infty} \frac{1}{(x+3)^3} dx = \int_4^{\infty} \frac{1}{u^3} du$ .
- c) The integral diverges. By comparison with  $\int_1^{\infty} \frac{1}{x}$
- d) The integral diverges.  $\lim_{x \rightarrow \infty} \frac{x^5}{e^{-x} + 1} = \infty$ . In order for the improper integral to converge, the integrand must approach 0.

17. b) and c)

18. a) volume of slice  $\approx 16\sqrt{36 - (6-x)^2} \Delta x$  Using Pythagorean Theorem.

b) volume of solid  $\approx \sum 16\sqrt{36 - (6-x_i)^2} \Delta x$  Using the notation of the text.

c) volume =  $\int_0^6 16\sqrt{36 - (6-x)^2} dx$

19. a)  $\int_0^1 3\sqrt{x} dx + \int_1^2 (6-3x) dx$

b)  $\int_0^3 \left( \frac{6-y}{3} - \frac{y^2}{9} \right) dy$

20. a)  $75\pi - \int_0^3 \pi(5e^{-x})^2 dx = \frac{(125 + 25e^{-6})\pi}{2}$

b)  $\int_0^3 \pi(5 - 5e^{-x})^2 dx = \frac{(75 + 100e^{-3} - 25e^{-6})\pi}{2}$

21. a)  $\int_0^8 \pi(y^{1/3})^2 dy = \frac{96}{5}\pi$

b)  $\int_0^8 \pi(y^{1/3} + 2)^2 dy - 32\pi = \frac{336}{5}\pi$

22.  $\int_0^2 (\pi(f(x))^2 - \pi(g(x))^2) dx + \int_2^7 (\pi(g(x))^2 - \pi(f(x))^2) dx$

23. a)  $\int_0^{\pi} (\sin x)^2 dx = \frac{\pi}{2}$

b)  $\int_0^{\pi} \frac{1}{2} \pi \left( \frac{\sin x}{2} \right)^2 dx = \frac{\pi^2}{16}$

24. Left hand rule:

$$10 \cdot \pi \left( \frac{26}{2\pi} \right)^2 + 10 \cdot \pi \left( \frac{22}{2\pi} \right)^2 + 10 \cdot \pi \left( \frac{18}{2\pi} \right)^2 + 10 \cdot \pi \left( \frac{12}{2\pi} \right)^2 + 10 \cdot \pi \left( \frac{6}{2\pi} \right)^2 = \frac{4160}{\pi} \text{ cubic inches}$$

Right hand rule:

$$10 \cdot \pi \left( \frac{22}{2\pi} \right)^2 + 10 \cdot \pi \left( \frac{18}{2\pi} \right)^2 + 10 \cdot \pi \left( \frac{12}{2\pi} \right)^2 + 10 \cdot \pi \left( \frac{6}{2\pi} \right)^2 + 10 \cdot \pi \left( \frac{2}{2\pi} \right)^2 = \frac{2480}{\pi} \text{ cubic inches}$$

Trapezoid rule:  $\frac{3320}{\pi}$  cubic inches      Use the average of the left and right hand rules.

$$25. \text{ a) } \int_0^1 \left( \pi (y^3)^2 - \pi (\sqrt{y})^2 \right) dy \qquad \text{b) } \int_0^\infty \pi \left( \frac{1}{x^2+1} \right)^2 dx$$

$$26. \int_0^5 (2 + 0.5 \cosh x) dx = 10 + 0.5 \sinh 5 \text{ grams}$$

$$27. \text{ a) } \int_0^8 \delta(x) \cdot 2\pi x dx \qquad \text{b) } \int_{-8}^8 \delta(x) \cdot 2\sqrt{64-x^2} dx$$

$$28. \int_0^{25} \left( -\frac{8}{5}h + 90 \right) \pi (10)^2 dh = 175,000\pi \text{ pounds}$$

$$29. \text{ a) } a_n = \frac{(-1)^n 2n}{(n+2)^2} \qquad \text{b) } \lim_{n \rightarrow \infty} a_n = -\frac{3}{5} \qquad \lim_{n \rightarrow \infty} b_n = 5$$

$$30. \text{ a) } \frac{(3/64)(1-(1/4)^8)}{1-1/4} = \frac{65535}{1048576} \qquad \text{b) } \frac{3/4}{1-1/4} = 1$$

$$31. P_n = \frac{(0.05)(200)(1-0.05^{n-1})}{1-0.05} \qquad Q_n = \frac{(200)(1-0.05^n)}{1-0.05} \qquad n = 1, 2, 3, \dots$$

32. a) The series converges.  $\int_2^{\infty} \frac{1}{x(\ln x)^2} dx = \frac{1}{\ln 2}$  Use the method of section 7.7 to evaluate the improper integral.

b) The series diverges.  $\int_1^{\infty} \frac{3x^2 + 2x}{\sqrt{x^3 + x^2 + 1}} dx = \infty$  Use the method of section 7.7 to evaluate the improper integral.

33. a) The series diverges.  $\lim_{n \rightarrow \infty} \frac{e(n)^2}{2(n+1)^2} = \frac{e}{2} > 1$

b) The series converges.  $\lim_{n \rightarrow \infty} \frac{(n+1)^2}{(2n+2)(2n+1)} = \frac{1}{4} < 1$

34. The Ratio Test gives us 1 and does not tell us anything about the convergence or divergence of the series.

~~35. The given series is conditionally convergent.  $\sum_{k=5}^{\infty} \frac{(-1)^{k-1}}{k(\ln k)}$  converges by the Alternating Series Test, while  $\sum_{k=5}^{\infty} \left| \frac{(-1)^{k-1}}{k(\ln k)} \right|$  diverges by the Integral Test.~~

36. a) and b)

37. a) False b) True c) False d) True e) False

38. a) The radius of convergence is  $R = 3$ . The interval of convergence is  $(-7, -1)$ .

b) The radius of convergence is  $R = \infty$ . The interval of convergence is  $(-\infty, \infty)$ .

c) The radius of convergence is  $R = 0$ . The series only converges for  $x = 1$ .

39. a) True b) True c) Impossible to determine.

40.  $P_2(x) = 4 + \frac{1}{3}(x-1) - \frac{1}{144}(x-1)^2$      $f(2) \approx P_2(2) = \frac{623}{144} \approx 4.3264$

41. The sign of  $c_0$  cannot be determined,  $c_1 > 0$ ,  $c_2 < 0$ .

42. a)  $f(3) = -1$ .      b)  $f'(3) = \frac{1}{2}$ .      c)  $f''(3) = -\frac{1}{6}$ .

d)  $\sum_{k=0}^{\infty} (-1)^{k+1} \frac{k!}{(2k)!} 3^k (x-1)^k$     Substitute  $3x$  into the series, then simplify.

43.  $\frac{1}{12}$       Integrate term by term. The result is recognizable as the series for  $\frac{x}{1-x}$ .

44.  $\cos(2\theta) = -\frac{1}{2} - \sqrt{3}\left(\theta - \frac{\pi}{3}\right) + \left(\theta - \frac{\pi}{3}\right)^2 + \frac{2\sqrt{3}}{3}\left(\theta - \frac{\pi}{3}\right)^3 + \dots$

45. a)  $\sin 1$     b)  $\ln(1.5)$     c) Series diverges because  $\frac{\pi}{e} > 1$ .

46.  $-\frac{1}{11}$       Use the series for  $\sin x$  to find the series for  $\frac{\sin x}{x}$ , then consider the term containing  $x^{11}$ .

47. a)  $x \ln(1+2x) = 2x^2 - \frac{4x^3}{2} + \frac{8x^4}{3} - \frac{16x^5}{4} + \dots = \sum_{n=0}^{\infty} \frac{(-1)^n 2^{n+1} x^{n+2}}{n+1}$

b)  $e^{-x^2} = 1 - x^2 + \frac{x^4}{2!} - \frac{x^6}{3!} + \dots = \sum_{n=0}^{\infty} \frac{(-1)^n x^{2n}}{n!}$

48.  $\frac{a}{(a+r)^2} = \frac{1}{a} \left(1 + \frac{r}{a}\right)^{-2} = \frac{1}{a} \left(1 - 2\left(\frac{r}{a}\right) + 3\left(\frac{r}{a}\right)^2 - 4\left(\frac{r}{a}\right)^3 + \dots\right)$

49. a) False    b) False    c) False

$$50. f(x) = \int_0^x \tan^{-1}(t) dt = \int_0^x \left( t - \frac{t^3}{3} + \frac{t^5}{5} - \frac{t^7}{7} + \dots \right) dt = \frac{x^2}{2} - \frac{x^4}{12} + \frac{x^6}{30} - \frac{x^8}{56} + \dots$$

The Taylor series for  $\tan^{-1}(t)$  at  $t = 0$  would be given.

51.  $Q = 0$  stable,  $Q = 1$  unstable

53. a) *ii*      b) *iii*      c) *i*      d) *iv*

$$54. y(t) = e^{3/4} e^{-(1/4)t} \quad \text{or} \quad y(t) = e$$

$$55. \text{ a) } y(x) = \frac{1}{2} x \sqrt{4-x^2} + 2 \arcsin\left(\frac{x}{2}\right) - 1 - \frac{\sqrt{3}}{2} - \frac{\pi}{3}$$

$$\text{ b) } x(\theta) = \frac{1}{2} \sin \theta \cos \theta + \frac{\theta}{2} + 1 - \frac{\pi}{2}$$

$$\text{ c) } y(t) = -\frac{3}{2} e^{4t^2} + \frac{1}{2} \quad \text{Watch for the sign issues when you remove the absolute values.}$$

$$\text{ d) } y(x) = 2 \sin\left(x + \frac{\pi}{6}\right)$$

$$56. \text{ a) } \frac{dQ}{dt} = -\alpha Q \text{ where } \alpha > 0 \quad \text{ b) } Q(t) = A e^{-\alpha t} \quad \text{ c) } t = \frac{7 \ln(90)}{\ln(9/5)} \approx 53.59 \text{ hours}$$

57. a) *ii*      b) *iii*      c) *i*      d) *iv*      e) *vi*      f) *v*

$$58. \text{ a) } \frac{dL}{dt} = 4 - 0.6L, \quad L(t) = \frac{A e^{-0.6t} + 4}{0.6}$$

b) The stable equilibrium solution is  $L = \frac{20}{3}$ . If we start with  $20/3$  grams per square centimeter of leaves, we will always have that amount.

59. a) *i*      b) *ii*      c) *iii*      d) *iv*



60. a)  $r(t) = \frac{1}{15}t + 60$

b)  $\frac{dH}{dt} = k\left(H - \left(\frac{1}{15}t + 60\right)\right)$   $H(0) = 180$  where  $k < 0$

61.  $\frac{dA}{dt} = 0.06\sqrt{A}$ ,  $A(t) = (0.03t + c)^2$

62. a)  $\frac{dP}{dt} = k(M - P)$  where  $k > 0$

b)  $\frac{dP}{dt} = kP(M - P)$  where  $k > 0$



63. a)  $\int_0^{15} 62.4(h)\pi\left(\frac{8}{15}h\right)^2 dh$

b)  $\int_0^{15} 62.4(h+3)\pi\left(\frac{8}{15}h\right)^2 dh$

c)  $\int_0^{10} 62.4(h)\pi\left(\frac{8}{15}h\right)^2 dh$

d)  $\int_3^{15} 62.4(h+3)\pi\left(\frac{8}{15}h\right)^2 dh$

64.  $500 \cdot 45 + \int_0^{45} 3(45 - x)dx = 25,537.5$  foot-pounds

One possibility.

65.  $\int_0^{10} 9.8\left(\frac{8}{0.5 \cdot 24 \cdot 10}\right)(10 - h)\left(\frac{12}{5}h\right)dh$  Joules

One possibility.

~~66.  $\int_0^{40} 62.4(40 - h)200dh$  pounds One possibility.~~